

## Extraction and purification of biosurfactant by locally isolated *Enterococcus faecium* bacteria and study of its antimicrobial activity

Dijlah A. Alimeer<sup>1</sup>, Nibras N. Mahmood<sup>2</sup> and Mohammad M. F. Al-Halbosi<sup>2</sup>

<sup>1,2</sup>Department of Biology, College of Science, Mustansiriyah University, Baghdad, Iraq;<sup>2</sup> Biotechnology Research Center/  
Al-Nahrain University/Iraq

\*Corresponding author's e-mail: [dijlamajeed@gmail.com](mailto:dijlamajeed@gmail.com)

In this research, the biosurfactant production by *Enterococcus faecium* isolated from Raw cow's milk was studied when growing in the MRS medium containing Lactose as a carbon source without Tween-80. *E. faecium* supernatant exhibited emulsion action of 60% towards kerosene oil as oil in water, while the surface activity was 34 mm. Biosurfactants with surface tension activity properties were extracted and partially purified from *E. faecium* by cold acetone 3:1. The chemistry structure of the biosurfactant was analyzed, which indicated that it is a glycol-lipo-peptide. Biosurfactant that purified from *E. faecium* showed antibacterial activity by reducing growth of several pathogenic G-ve and G+ve bacteria, including *aureus*, *Staphylococcus epidermidis*, *Streptococcus* sp., *Staphylococcus Escherichia coli*, *Klebsiella* sp., *Pseudomonas aeruginosa*, *Acinetobacter* sp. and *Serratia* sp. The inhibitory activity of biosurfactant against pathogenic bacteria was studied by using agar diffusion method, where the biosurfactant had higher inhibitor activity against *E. coli*, where highest diameter of inhibition zone was (23mm). At all concentrations examined, a biosurfactant derived from *E. faecium* considerably decreased bacterial adhesion, albeit to a lower amount.

**Keyword:** *Enterococcus faecium*, LAB, Biosurfactant, Antibacterial and Antiadhesive activity.

### INTRODUCTION

Biosurfactants are hydrophobic and hydrophilic amphiphilic compounds constructed by a category of microorganisms, including bacteria, fungus, and yeast. These molecules display reduced surface tension or interface tension as well as emulsification properties. They are either synthesized extracellularly or bound to cell surfaces (Eras-Munoz *et al.*, 2022). In contrast to chemical surfactants, biosurfactants have many benefits, including broad chemical variety, low toxicity, strong biocompatibility, high selectivity, good environmental compatibility, and high biodegradability. (Inamuddin *et al.*, 2019; Rodrigues 2015). The advantage of biosurfactant is that they can act in conditions of high pH, temperature, and salt content. It can be produced on several types according to the microorganism produced, such as: lipopeptides, Glycolipid and Glycolipoprotein are showed encouraging antibacterial, antifungal, antiviral, and antiadhesive properties (Moldes *et al.*, 2021). Which they can be used instead of traditional antibiotics to combat different food-borne pathogens. There have been reports of different kinds of biosurfactants being produced by bacterial genera including, *Pseudomonas*,

*Bacillus*, *Acinetobacter*, *Halomonas*, *Arthrobacter*, *Lactobacillus* and *Enterococcus* (Giri *et al.*, 2019). The goal of this study was to explore the antibacterial and anti-adhesive characteristics of biosurfactant supplied By *Enterococcus faecium* against several pathogenic bacteria Collected from the educational laboratories of the Department of Life Sciences at Al-Mustansiriyah University.

### MATERIALS AND METHODS

**Microorganisms:** *Enterococcus faecium* was isolated from Row cow's milk samples taken from Al-Fadilia cow farm., Baghdad Iraq, samples were transferred to the lab at 4°C. An aliquot (1 gm) was homogenized in 9 mL of sterile 0.9% (w/v) NaCl, diluted 10-fold, and plated on Man Rogosa Sharp agar (MRS) After development the inoculum from the culture flasks was subcultured on bile Esculin agar as confirmative of *Enterococcus* spp. then subcultured on the MRS agar to obtain pure colonies. These isolates were stored at -20°C in MRS broth containing 20% glycerol stock till they were used in this study. Then identified throughout cultural, microscopical, biochemical test and Vitek-2 according to

Alimeer, D.A., N.N. Mahmood and M.M. F. Al-Halbosi. 2023. Extraction and purification of biosurfactant by locally isolated *Enterococcus faecium* bacteria and study of its antimicrobial activity. Journal of Global Innovations in Agricultural Sciences 11:543-548.

[Received 23 July 2023; Accepted 15 Aug 2023; Published 22 Dec 2023]



Attribution 4.0 International (CC BY 4.0)

(Teixeira *et al.*, 2015). Biosurfactant produced from *Enterococcus faecium* was subjected to the test against eight isolates, including *Staphylococcus epidermidis*, *Streptococcus pyogenes*, *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, and *Serratia marcescens*. These isolates were obtained from various clinical samples of Iraqi patient, subsequently identified by cultural, microscopic, and biochemical tests in accordance with the guidelines laid out by Forbes (2002) and confirm it with Vitek-2.

**Cultivation of *E. faecium* and biosurfactant Production:** The MRS broth utilized for Biosurfactant producing *E. faecium* isolation contains (20 g D-Lactose l<sup>-1</sup>, 10 g peptone l<sup>-1</sup>, 10 g Beef extract l<sup>-1</sup>, 5 g Yeast extract l<sup>-1</sup>, 2g Dipotassium phosphate l<sup>-1</sup>, 2g Tri-ammonium hydrogen citrate l<sup>-1</sup>, 5g sodium acetate l<sup>-1</sup>, 0.2g Magnesium sulphate l<sup>-1</sup> and Manganese sulfate l<sup>-1</sup> without Tween-80. For biosurfactant production, 10 ml MRS broth was inoculated with a fresh culture of *E. faecium* and developed at 37°C for 48hrs. and further used as the seed culture for production of biosurfactant. (Salman and Alimer 2014).

**screening for glycolipoprotein biosurfactant production Oil spreading method:** In order to perform the oil-spreading test, water and oil were placed in a petri dish. A drop of supernatant and put on the oil's surface. For each isolate, the diameter of the obvious zone on the oil surface was measured three times. PBS was utilized as a negative controller (Morikawa *et al.* 2000).

**Determination of emulsification Index (E24%):** The Emulsification indicator test was done by following to Sharma *et al.* (2014) approach for assessing biosurfactant activity: The capacity of isolates to emulsify was evaluated using an emulsification index (E24) intended for kerosene oil. In a test tube, 5 mL of Kerosene was mixed with 5 mL of supernatant. The test tube was then rapidly vortexed for two minutes before being let to stand for 24 hours. The ratio of the emulsification index was anticipated using the equation which showed encouraging properties of antibacterial, antifungal, antiviral, and anti-adhesive.

$$E24\% = \frac{\text{Height of the emulsion layer}}{\text{Total height of the mixture}} \times 100$$

**Biosurfactant production, extraction, and partial purification:** For crude biosurfactant production in flask condition, a pre-culture of 2% (v/v) *E. faecium* was added to 1200 ml of MRS-Lactose (glucose was substituted via lactose) potage (pH 7) and cultivated for 48 hours at 37°C. Cells were separated by centrifugation after 48 hours of incubation (10,000 g, 10 min, 4°C). After cooling centrifuge, the supernatant containing biosurfactant was treated with cold acetone solvent (3:1) (v/v), The mix was surprised continuously at 200 rpm, 4 °C for 15-20 hours inside an incubator shaker, and then centrifuged at 10000 rpm for 15 minutes at 4 °C. The supernatant was discarded, the precipitate was collected and This served as partially purified

biosurfactant. The pellet was dissolved in 5 ml D.W., and pH was regulated at 7 with NaOH (1N) as prescribed by (Majid Hussein *et al.*, 2015).

**Estimation of Biomass:** After centrifugation in the biosurfactant extraction process, the cell pellet was cleaned, re-suspended in sterilized water, and centrifuged repeatedly. A cell pellet was later dehydrated until it attained a constant weight in an electric oven at 105°C (Schelegueda *et al.*, 2015). **chemical analysis of biosurfactant:** Lipid, carbohydrate and protein content were evaluated with Anschau (2017) method, phenol, and sulphuric acid method by (Dubois *et al.*, 1956) and Bradford method (Bradford, 1976), respectively.

#### Anti-bacterial Assay

**Agar diffusion method:** Pathogenic bacterial strains *Staphylococcus aureus*, *Escherichia coli*, *Streptococcus pyogenes*, *Klebsiella*, *Staphylococcus epidermidis*, *pneumonia*, *Pseudomonas aeruginosa*, *Acinetobacter baumannii* and *Serratia marcescens*, were obtained from various clinical samples of Iraqi patients and Muller-Hinton Agar (MHA) prepared plates were swapped with these pathogenic bacteria. Wells were made in the agar plates using a sterile well maker and then 100 µl of a biosurfactant with a 40 mg/ml concentration were added to a separate well. The plates were incubated for 24 hours at 37°C. The diameter of the clear zone was observed (Giri *et al.*, 2019 and Hussein and Hasan., 2022).

**Anti adhesion Assay:** According to the method established by Ali (2012), co-incubation studies were used to measure the anti-adhesion activity of the partially purified biosurfactant obtained from *E. faecium* against *Staphylococcus epidermidis*, *Staphylococcus aureus*, and *Streptococcus pyogenes*, *Escherichia coli*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa*, *Acinetobacter baumannii* and *Serratia marcescens*. The inhibition of adhesion ratios of partial purified biosurfactant for each pathogenic bacteria was determined as equation termed by Gudina *et al.* (2010).

$$\text{Inhibition of Adhesion \%} = \left[ 1 - \left( \frac{A}{A_0} \right) \right] \times 100$$

A: represent the absorbance of the well with a biosurfactant and A0: the absorbance of the control well.

## RESULTS AND DISCUSSION

**Initial isolation:** The bacterial isolate was examined based on its morphological and biochemical characteristics, All isolates grew on bile esculin agar. This media a choosy differential agar used to isolate and identify members of the genus *Enterococcus*. Esculin is the differentiating element, whereas bile salts are the selective ingredient. Esculin is hydrolyzed by *Enterococcus* into combined products with ferric citrate in the medium to form insoluble iron ions, Black hallow colonies on Bile Esculin agar was considered as presumptive enterococci colonies (Davis *et al.*, 2022). Enterococci were observed small, smooth, entire margin and white colonies on



MRS agar (Ben Braiek *et al.*,2017). and Identification was performed based on the Vitek2 system (Forbes *et al.*,2007; Sarwar *et al.*,2021).

**Screening for biosurfactant production:** In the current investigation, *E. faecium* isolates were screened for biosurfactant production by two methods: Oil spreading method and Emulsification Index 24%


**Oil Spreading test:** The tested *Enterococcus faecium* isolate showed the highest oil spreading activity with diameter 34 mm. Results are given in Fig. 1.

The results indicated the *E. faecium* had the ability to synthesize biosurfactants. The diameter of the clearance zone formed by biosurfactant-comprising solution has been shown to be directly proportional to the concentration of biosurfactant (Youssef *et al.*, 2005), Ghazi Faisal *et al.* (2023) reported that this method is better than other screening methods due to be rapid and easy to carry out, requires no specialized equipment. while Lamia *et al.* (2021) indicated that the development of clear zone on surface of oil is a unique quality of surfactants, which, together with tensile surface, provides an index to estimate the performance of biosurfactant-making bacteria .



**Figure 1. The spreading of partial biosurfactant produced by *E. faecium* on oil surface.**

**Emulsification activity E24%:** Emulsification index (E24%) of *E. faecium* was determined with bacterial supernatant and kerosene oil. *E. faecium* showed good emulsifying activity, emulsion formed with E24 (60%), Figure 2 shows the E24% result. According to Das *et al.* (2009) the emulsification index corresponds to the surfactant concentration. For another studies Lactic acid bacteria formed good emulsifying ability (Brzozowski *et al.*, 2011), While Ghasemi *et al.* (2019) reported that the Biosurfactant produced by lactic acid bacterium *Pediococcus dextrinicus* SHU1593 is able to emulsify n-hexadecane (58%) and kerosene (56%).



**Figure 2. The Emulsion form of *E. faecium***

A: phosphate buffer saline, B: *E. faecium* supernatant

#### **Extraction and Partial Purification of the Biosurfactants:**

The crude Biosurfactant produced by *E. faecium* was extracted and partially purified by using cold acetone: Supernatant cell culture (3:1) ( Al-Rrubaie *et al.*,2019). Arsyah *et al.*, (2018) also selected the cold acetone method for the extraction and purification of biosurfactants.

This occurs as a result of the availability of the hydrophobic end of the biosurfactant, which makes them soluble in acetone or organic solvent. (Shah *et al.*, 2016). In current study, the yield of partially purified biosurfactant was (9.4 g/2.1 gm Biomass /1 liter of culture medium) from *E. faecium* in the presence of Lactose as sole source of carbon.

The results of chemical composition analysis, the biosurfactant produced by *Enterococcus faecium* was content of Glycolipoprotein. This study indicates that the lipid content of purified Biosurfactant was 39.7% according to Anschau standard curve. According to Dubois standard curve the carbohydrate contents were 33% and the concentration of the protein was 9.5%, according to an analysis of the protein's absorbance using the Bradford standard curve.

**Antibacterial activity of biosurfactant:** Glycolipoprotein from *E. faecium*, showed antibacterial activity against tested isolates with different degree (Table 1). From those results, the Biosurfactant gave the largest inhibition zone (23 mm) against *E. coli*. Biosurfactant very efficiently suppressed the growth of tested G-negative and G- positive bacteria. It is significant to mention that the biosurfactant produced by *E. faecium* demonstrated good effect against bacterial infections. Various biosurfactants with antibacterial activity have already



been reported. However, few researchers have investigated the antibacterial action of biosurfactants isolated from *Enterococci*. Similarly, biosurfactant formed from *Lactobacillus rhamnosus* exhibited antimicrobial effect against some bacteria causing UTI in Iraqi women (Salman and Alimer; 2014), *Lactobacillus casei*-derived biosurfactant displayed antimicrobial activities against *S. aureus* strains (Merghni *et al.*, 2015). also, biosurfactant produced from *Bacillus* strain had the highest inhibitory zone against tested *E. coli* MTCC ( $13 \pm 0.5$  mm)(Giri *et al.*,2019), biosurfactant produced by *Pediococcus dextrinicus* had significant a complete antimicrobial effect against *E. coli*, *E. aerogenes*, and *P. aeruginosa* at concentration 25 mg/ml (Ghasemi *et al.*,2019), another studies, biosurfactant produced by Lactic Acid Bacteria appeared antibacterial result against Gram positive and Gram negative bacteria (Hassan, (2018) ; Dong *et al.*, 2019 and Nataraj *et al.*, 2021).

**Table 1. Antibacterial effect of biosurfactant produced by *E. faecium*.**

Pathogenic bacteria	Inhibition zones(mm) at 25 mg/ml
<i>K. pneumonia</i>	17
<i>S. epidermidis</i> ,	20
<i>S. pyogenes</i> ,	19
<i>P. aeruginosa</i>	15
<i>E. coli</i>	23
<i>A. baumannii</i>	12
<i>S. marcescens</i> .	16
<i>S. aureus</i>	21

The biosurfactant's mechanism of antibacterial activity focuses on the fact that the structural likeness to detergent provided by the existence of the hydrophilic head and lipophilic tail allows for its penetration into the lipid bilayer of the cell membrane, producing disruption (Malakar, and Deka, 2021). Another reasoning for how biosurfactants work as antimicrobials is the intercalation of biosurfactant into the membrane causes the formation of holes in the cell membrane. Also Nucleic acid leakage among other intercellular materials, other routes for the manifestation of antibacterial activities include interactions with membrane phospholipids or modifications to the electrical conductivity of membranes (Abdelli *et al.*,2019; Sharma and Sahran 2016). The other hand, the anti-adhesive effects of partial purified biosurfactant was assessed against tested pathogenic bacteria (Table 2). The partial purified Biosurfactant showed anti-adhesive activity against all tested bacteria, but anti-adhesive activity was dependent on biosurfactant concentrations the highest anti-adhesive percentage was observed for *E. coli* (78%) at a concentration of biosurfactant 50 mg/ml .

Biosurfactant produced from *E. faecium* against *E. coli* and *S. aureus* was comparable to the results attained with biosurfactant produced by *Lactobacillus brevis* and *Bacillus* sp. which inhibited adhesive of *E. coli* and *S.aureus* (Haddaji

*et al.*,2022). Shawkat *et al.* (2019) showed that the biosurfactant even at low to moderate concentrations, it proved efficient against *Salmonella enteritidis*, *Staphylococcus epidermis*, and *E. coli* but, low activity was demonstrated for *Staphylococcus aureus*. According to Rodrigues *et al.* (2006), the primary objective of a biosurfactant is to alter the surface's physicochemical characteristics in order to decrease the attraction that microorganisms have to its surface.

**Table 2. Anti-adhesive activity of partial purified biosurfactant produced by *E. faecium*.**

Pathogenic bacteria	Anti-adhesive percentage (%) of biosurfactant at mg/ml				
	3.12	6.25	12.5	25	50
<i>K. pneumonia</i>	4	5	5	7	10
<i>S. epidermidis</i> ,	10	18	27	43	59
<i>S. pyogenes</i> ,	12	16	21	39	64
<i>P. aeruginosa</i>	15	20	23	46	67
<i>E. coli</i>	13	19	20	43	78
<i>A. baumannii</i>	19	23	25	38	55
<i>S. marcescens</i> .	2	5	6	10	13
<i>S. aureus</i>	20	28	35	55	72

When compared to the control, positive percentages show a decrease in bacterial adhesion, the percentage 0% was selected as the negative control in order to demonstrate that the biosurfactant isn't present.

**Conclusion:** In conclusion, It was observed that *E.faecium* have the ability to create biosurfactant with antibacterial and antiadhesive property against some tested pathogenic isolates *Staphylococcus aureus* *Escherichia coli*, *streptococcus sp*, *Staphylococcus epidermidis*, *Klebsiella sp.*,*Pseudomonas aeruginosa*, *Acinetobacter sp.* and *Serratia sp*.

**Authors' contributions:** Dijlah A. Alimeer: the idea, procedure, writing and editing. Nibras N. Mahmood: Results analysis and editing. Mohammad M. F. Al-Halbosi : contributes to fieldwork and linguistic review.

**Funding:** ourselves

**Ethical statement:** Following the university of Al-Mustansiriyah ethical rules.

**Acknowledgement:** we thank all the staff of the Department of Biology, College of Science, Mustansiriyah University and Biotechnology Research Center/ Al-Nahrain University/Iraq.

**Code Availability:** Not applicable

**Consent to participate:** All agree to participate.





**Consent for publication:** All agree for publication. Abd El-Ghani A.A. 2004. Influence of diet supplementation with yeast culture (*Saccharomyces cerevisiae*) on performance of Zaraibi goats. *Small Ruminant Research* 52:223-229.

## REFERENCES

- Abdelli, F., M. Jardak., J. Elloumi, D. Stien, S. Cherif, S. Mnif and S. Aifa. 2019. Antibacterial, anti-adherent and cytotoxic activities of surfactin(s) from a lipolytic strain *Bacillus safensis* F4. *Biodegradation* 30:287-300.
- Ali, O.A. 2012. Prevention of *Proteus mirabilis* Biofilm by Surfactant Solution. *Egyptian Academic Journal of Biological Sciences* 4:1- 8.
- Al-Rubaie, G., N. H. Zaki, and S. Latif. 2019. Antimicrobial Activity of Freshwater *Cyanobacterium* Westielloopsis prolifica. *Al-Mustansiriyah Journal of Science* 29:42-49.
- Anschau, A., C. S. Caruso, R. C. Kuhn and T. T. Franco. 2017. Validation of the sulfo-phospho-vanillin (SPV) method for the determination of lipid content in oleaginous microorganisms. *Brazilian Journal of Chemical Engineering* 34:19-27.
- Arsyah, D.M., E. Kardena, and Q. Helmy. 2018. Characterization of biosurfactant produced by petrofilic bacteria isolated from hydrocarbon impacted soil and its potential application in bioremediation. In *IOP Conference Series: Earth and Environmental Science* 106:012101.
- Ben Braiek, O., H. Ghomrassi, P. Cremonesi, S. Morandi, Y. Fleury, P. Le Chevalier, K. Hani, O. Hadj and T. Ghrairi. 2017. Isolation and characterisation of an enterocin P-producing *Enterococcus lactis* strain from a fresh shrimp (*Penaeus vannamei*). *Antonie van Leeuwenhoek* 110:771-786.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry* 72:248-254.
- Brzozowski, B., W. Bednarski and P. Gol'ek. 2011. The Adhesive Capability of Two *Lactobacillus* Strains and Physicochemical Properties of Their Synthesized Biosurfactants. *Food Technology and Biotechnology* 49 :177-186.
- Das, P., S. Mukherjee and R. Sen. 2009. Substrate dependent production of extracellular biosurfactant by a marine bacterium. *Bioresource Technology* 100:1015-1019.
- Davis, B. C., I. Keenum, J. Calarco, K. Liguori, E. Milligan, A. Pruden and V. J. Harwood. 2022. Towards the standardization of *Enterococcus* culture methods for waterborne antibiotic resistance monitoring: A critical review of trends across studies. *Water Research X*, 100161.
- Dong, Y., G. Shu, C. Dai, M. Zhang and H. Wan. 2019. Screening and identification of biosurfactant-producing lactic acid bacteria. *Acta Universitatis Cibiniensis. Series E: Food Technology* 23:85-92.
- DuBois, M., K.A. Gilles, J.K.Hamilton, P.T. Rebers and F. Smith. 1956. Colorimetric method for determination of sugars and related substances. *Analytical chemistry* 28:350-356.
- Eras-Munoz, E., A. Farré, A. Sánchez, X. Font and T. Gea. 2022. Microbial biosurfactants: a review of recent environmental applications. *Bioengineered* 13:12365-12391.
- Forbes, B.A., D.F. Sahm and Weissfeld 2007. *Bailey and Scott's Diagnostic microbiology* 12th. Edition St Louis, Mosby Elsevier, China.pp. 842-855.
- Forbes, B.A., Saham, D.F. and Weissfeld, A.S. 2002. *Diagnostic Microbiology*. 10th ed. Mosby. Inc. St. Louis, Mo.
- Ghasemi, A., M. Moosavi-Nasab, P. Setoodeh, G. Mesbahi and G.Yousefi. 2019. Biosurfactant production by lactic acid bacterium *Pediococcus dextrinicus* SHU1593 grown on different carbon sources: strain screening followed by product characterization. *Scientific reports* 9:5287.
- Ghazi Faisal, Z., M. Sallal Mahdi and K.H. Alobaidi 2023. Optimization and Chemical Characterization of Biosurfactant Produced from a Novel *Pseudomonas guguanensis* Strain Iraqi ZG. KM. *International Journal of Microbiology*, 2023.
- Giri, S.S., E.C. Ryu, V. Sukumaran and S.C. Park. 2019. Antioxidant, antibacterial, and anti-adhesive activities of biosurfactants isolated from *Bacillus strains*. *Microbial pathogenesis* 132:66-72.
- Gudina, E.J., V. Rocha, J.A. Teixeira and L.R. Rodrigues. 2010. Antimicrobial and antiadhesive properties of a biosurfactant isolated from *Lactobacillus paracasei ssp. paracasei* A20. *Letters in applied microbiology* 50:419-424.
- Haddaji, N., A. Mahdhi, N.Bouali, M. Ghorbel,, O. Bechambi, N. Leban, M. Ameer and R. Mzoughi. 2022. Biosurfactants as inhibitors of the adhesion of pathogenic bacteria. *Emirates Journal of Food and Agriculture* 34:36-43.
- Hassan, Z. 2018. The biosurfactant and antimicrobial activity of lactic acid bacteria isolated from different ources of fermented foods. *Asian Journal of Pharmaceutical Research and Development* 6:60-73.
- Hussein, L.M. and A.Y. Hasan. 2022. The Antibacterial Activity of the Biosynthesized Copper Oxide Nanoparticles by Lantana camara Flowers Extract Against Some Bacterial Isolated from Burns. *Al-Mustansiriyah Journal of Science* 33:39-52.
- Inamuddin, A. M., Asiri, A., & Suvardhan, K. (2019). *Green sustainable process for chemical and environmental engineering and science*. Amsterdam, The Netherlands: Elsevier.



- Karnwal, A., S. Shrivastava, A. R. M. S. Al-Tawaha, G. Kumar, R. Singh, A. Kumar, A. Mohan, Yogita and T. Malik. 2023. Microbial Biosurfactant as an Alternate to Chemical Surfactants for Application in Cosmetics Industries in Personal and Skin Care Products: A Critical Review. *BioMed Research International*, 2023.
- Lamia, K., E.A. Kamel, E. Sonia, M.I. Soudes and Z.I. Billal. 2021. Production of biosurfactant from a novel strain *Bacillus amyloliquefaciens* K2E. *Revue Agrobiologia* 11:2628-2640.
- Majid Hussein, A.J., M. Nasir and M. Aziz. 2015. Characterization of Biosurfactant Produced by a Novel Thermophilic Strain (*Geobacillus thermoleovorans* JQ 912239). *Asian Journal of Biotechnology* 7:96-107.
- Malakar, C. and S. Deka 2021. Biosurfactants Against Drug-Resistant Human and Plant Pathogens: Recent Advances. *Biosurfactants for a Sustainable Future: Production and Applications in the Environment and Biomedicine* 353-372.
- Merghni, A., M.B. Nejma, I. Helali, H. Hentati, A. Bongiovanni, F. Lafont, M. Aouni, M. Mastouri. 2015. Assessment of adhesion, invasion and cytotoxicity potential of oral *Staphylococcus aureus* strains. *Microbial pathogenesis* 86:1-9.
- Moldes, A.B., L. Rodríguez-López, M. Rincón-Fontán, A. López-Prieto, X. Vecino and J.M. Cruz. 2021. Synthetic and bio-derived surfactants versus microbial biosurfactants in the cosmetic industry: An overview. *International Journal of Molecular Sciences* 22:2371.
- Morikawa, M., Y. Hirata and T. Imanaka. 2000. A study on the structure–function relationship of lipopeptide biosurfactants. *Biochimica et Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids*, 1488: 211-218.
- Nataraj, B.H., C. Ramesh and R.H. Mallappa. 2021. Characterization of biosurfactants derived from probiotic lactic acid bacteria against methicillin-resistant and sensitive *Staphylococcus aureus* isolates. *LWT*. 151: 112195.
- Rodrigues, L. R. 2015. Microbial surfactants: fundamentals and applicability in the formulation of nano-sized drug delivery vectors. *Journal of colloid and interface science* 449:304-316.
- Rodrigues, L., I. M. Banat, J. Teixeira and R. Oliveira. 2006. Biosurfactants: potential applications in medicine. *Journal of antimicrobial chemotherapy* 57:609-618.
- Salman, J.A.S. and D.A. Alimer. 2014. Antibacterial and antiadhesive properties of a biosurfactant isolated from *Lactobacillus rhamnosus* against some bacteria causing UTI in Iraqi women. *International Journal of Current Research* 6:5368-5374.
- Sarwar, M., S. A. H. Gardezi, G. Zaman, A. Ikram, W. Hussain and M.T. Khadim. 2021. Diagnostic accuracy of in-house biochemical testing for identification of *Enterococcus* species isolated from various clinical specimens against Vitek 2 system. *Pakistan Armed Forces Medical Journal* 71:155-61.
- Schelegueda, L. I., M. Vallejo, M. F. Gliemmo, E. R. Marguet and C.A. Campos. 2015. Synergistic antimicrobial action and potential application for fish preservation of a bacteriocin produced by *Enterococcus mundtii* isolated from *Odontesthes platensis*. *LWT-Food Science and Technology* 64:794-801.
- Shah, A., S. Shahzad, A. Munir, M. N. Nadagouda, G. S. Khan, D. F. Shams, D.D. Dionysiou and U.A. Rana. 2016. Micelles as soil and water decontamination agents. *Chemical reviews* 116:6042-6074.
- Sharma M, D. R. Modi, M. Saxena. 2014. Role of Lactic Acid Bacteria as Probiotics in Health and Disease. *Prevention* 82:1-15.
- Sharma, D. and B.S. Saharan. 2016. Functional characterization of biomedical potential of biosurfactant produced by *Lactobacillus helveticus*. *Biotechnology Reports* 11:27-35.
- Shawkat, S. M., M. Al-Jawasim and Z. F. Hussain. 2019. Production of anti-adhesive coating agent for table eggs by *Lactobacillus acidophilus* MTCC 447. *Plant Arch* 19:1795-1798.
- Teixeira, L. M., M. D. G. S. Carvalho, R. R. Facklam and P. L. Shewmaker. 2015. *Enterococcus*. *Manual of clinical microbiology* 403-421.
- Youssef, N. H., K. E. Duncan and M. J. McInerney. 2005. Importance of 3-hydroxy fatty acid composition of lipopeptides for biosurfactant activity. *Applied and Environmental Microbiology* 71:7690-7695.

